

#18 p/anner
12.20.00


Docket No. 47653.1 (1789)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: J.C. Houck, et al.
Serial Number: 09/189,130 Art Unit: 1631
Filed: November 10, 1998 Examiner: M. Borin
For: SMALL PEPTIDES AND METHODS FOR TREATMENT OF
ASTHMA AND INFLAMMATION

CERTIFICATE OF MAILING

I hereby certify that this paper (along with any paper referred to as being attached or enclosed) is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner for Patents, Washington, D.C. 20231 on December 4, 2000.


Barbaranne Jenness

Honorable Commissioner for Patents
Washington, DC 20231

Sir:

DECLARATION OF JAMES CLAGETT

I, James Clagett, hereby declare that:

1. I am a citizen of the United States of America residing at 5615 139th Avenue SE, Snohomish, Washington 98290 and I am one of the Applicants in the above application.

2. I hold a Ph.D. in microbiology from the University of Nebraska. I have over 30 years experience in research and development related to microbiology, particularly in the fields of immunology and immunopathology. A copy of my curriculum vitae is

Declaration of James Clagett, Ph.D.
Ser. No. 09/189,130
Page 2 of 8

2

attached hereto as Attachment A.

3. Since 1997, I have been a consultant providing scientific expertise to the biotechnology and pharmaceutical communities.

4. I personally performed or directly oversaw the experiments which produced the results presented and discussed herein.

5. I have read and understand the Office Action of June 28, 2000, including the references cited therein.

6. The present invention is directed to a pharmaceutical composition having anti-inflammatory activity comprising a pharmacological carrier and an anti-inflammatory effective amount of a peptide having the formula f-Met-Leu-Phe-Phe.

7. Based on my knowledge and experience in the field, it is my opinion that, prior to the present invention, it was well known to those skilled in the art that formyl methionyl peptides have pro-inflammatory activity.

8. However, surprisingly, we have discovered that f-Met-Leu-Phe-Phe, can provide a useful anti-inflammatory effect.

9. In the Office Action dated June 28, 2000, the examiner states:

Declaration of James Clagett, Ph.D.
Ser. No. 09/189,130
Page 3 of 8

3

... the essential difference [is] the effect of a biological mediator (such as f-Met peptide) when it is used alone as compared to its use in the presence of another pro-inflammatory agent. Cellular response to f-Met peptides (which can be described as inflammatory response) is the same type of reaction which mediates response of the organism to a foreign infection. It is well known in the art that biological mediators such as chemotactic factors stimulate the migration of neutrophils from circulation into sites of infection or tissue damage. These mediators are also believed to increase cell adhesion to injured sites and to activate neutrophils to release toxic agents such as oxygen metabolites and proteases. Thus, in the presence of a provoked infection the response caused by f-Met peptides have protective, anti-inflammatory function.

10. It is true that the prior art teaches that:

Cellular response to f-Met peptides (which can be described as inflammatory response) is the same type of reaction which mediates response of the organism to a foreign infection. It is well known in the art that biological mediators such as chemotactic factors stimulate the migration of neutrophils from circulation into sites of infection or tissue damage. These mediators are also believed to increase cell adhesion to injured sites and to activate neutrophils to release toxic agents such as oxygen metabolites and proteases

However, those responses are "pro-inflammatory" responses. The claimed composition of the present invention blocks those responses. Thus, the claimed composition has an "anti-inflammatory" response.

11. The examiner also states and concludes that:

Characteristically, ... the effect of the claimed composition is demonstrated only as inhibitor of inflammatory effect caused by another f-Met peptide, fMLP. The absence ... of showing of the effect of fMLPP alone is not surprising because Kermode shows (Table 2) that fMLPP (the peptide of the claimed composition) is more potent chemotactic agent and stimulator of neutrophil degranulation than fMLP (the peptide used as "pro-inflammatory" agent). One would expect that fMLPP, alone, would be at least as "pro-inflammatory" as fMLP.

Declaration of James Clagett, Ph.D.
Ser. No. 09/189,130
Page 4 of 8

4

12. However, Kermode conducted *in vitro* tests using rabbit neutrophils that were suspended in solution containing salts, BSA, buffer and for some tests glucose. We have found that such *in vitro* tests are not a predictor of the bioactivity of fMLPP *in vivo*. Although based on the teachings of the prior art, " [o]ne would expect that fMLPP, alone, would be at least as 'pro-inflammatory' as fMLP," as concluded by the examiner, that is an erroneous expectation. Further, based on the teachings of the prior art, one of ordinary skill in the art would not expect fMLPP to act any differently after prior treatment with fMLP.

13. To satisfy the examiner's curiosity about the effect of fMLPP alone, the following experiments have been performed under my direction and control.

Briefly, (i) 200 μ g of fMLP alone; (ii) 200 μ g fMLPP (HK-X) alone; (iii) 200 μ g of fMLP and 200 μ g of HK-X administered simultaneously; and (iv) as a control the vehicle of HK-X (4% DMSO in Tyrode's solution) each were injected subcutaneously into the dorsum of the feet of female Balb/CJ female mice. Animals were sacrificed at 30 minutes post-injection and the feet collected for histological examination. The cutaneous soft tissues were dissected from the bones, fixed in 10% (v/v) neutral buffered saline and embedded in paraffin. Five - seven micron sections were cut and stained with H&E for the detection of cellular content and location within the muscularis and dermis. The numbers of neutrophils in the extravascular zone, occupying 2,200 μ^2 around blood vessels, were counted. For each specimen, a total of 10 vessels were counted. The means and SE are presented. Differences between groups were determined by one-way ANOVA followed by the appropriate post hoc tests

Declaration of James Clagett, Ph.D.
Ser. No. 09/189,130
Page 5 of 8

5

using SigmaStat version 2.0 software (SPSS Inc., Chicago, IL). The results are illustrated in the figures attached hereto.

14. Figure 1 shows the polymorphonuclear cell count outside the vessels, 30 minutes after injection of 200 μ g fMLP alone; 200 μ g fMLP and 200 μ g HK-X together; 200 μ g HK-X alone; and vehicle alone, into the subcutaneous tissues of the dorsum of mice feet.

- Injection of 200 μ g of fMLP alone resulted in 24 neutrophils per 2,200 μ^2 in the intercellular matrix.
- In specimens from treatment with HK-X alone, results were substantially identical to the control (vehicle treatment alone) ($p < 0.05$).
- The simultaneous exposure of fMLP and HK-X resulted in a 91 % inhibition of neutrophils diapedesis compared to fMLP alone ($p < 0.05$).

15. Figure 2 is a microphotograph comparing stained tissue sections harvested from mice 30 minutes after injection with (i) 200 μ g of fMLP alone, (ii) 200 μ g fMLP and 200 μ g HK-X simultaneously and (iii) 200 μ g of HK-X alone into the subcutaneous layers of the skin on the dorsum of mice feet.

- Panel A shows the results of injection of 200 μ g of fMLP alone. FMLP treated mouse skin shows the influx of the neutrophils in the surrounding areas of the blood vessel (BV). The neutrophils are located in the connective tissue area (Arrows).

Declaration of James Clagett, Ph.D.
Ser. No. 09/189,130
Page 6 of 8

6

- Panel B shows the results of 200 µg fMLP and 200 µg HK-X when injected simultaneously. The mouse skin shows no cellular infiltration at the surrounding areas of the blood vessel (BV). Occasionally, neutrophils are seen. (Arrows).
- Panel C shows the results of injection of 200 µg of HK-X alone. The treated mouse skin shows no changes in connective tissues and the surrounding area of the blood vessel (BV).

16. Figure 3 shows a higher power examination of specimen similar to those in figure 2.

- Panel A shows the subcutaneous tissue of an HK-X treated mouse. The subcutaneous tissue is normal. The blood vessels (BV) show no neutrophils. Only occasionally a neutrophil is observed (Arrow).
- Panel B is the vehicle control treated mouse. The subcutaneous tissue shows no sign of the effect of a vehicle solution injection.
- Panel C shows the results of 200 µg fMLP and 200 µg HK-X when injected simultaneously. The subcutaneous tissue shows very little change in the histology except for a few neutrophils attached on the endothelial cell surface (Arrow heads). Neutrophils are observed but in very little numbers as compared to the fMLP treated skin tissue (Panel D).
- Panel D shows the results of shows the results of injection of 200 µg fMLP alone. The subcutaneous tissue was heavily infiltrated with

Declaration of James Clagett, Ph.D.
Ser. No. 09/189,130
Page 7 of 8

7

neutrophils 30 minutes after injection of fMLP as seen in the surrounding area of the blood vessel (BV). Many neutrophils are also seen attached to the surface of endothelial cells.

17. Based on my knowledge and experience in the art, from these experiments, it can be concluded that fMLP (prior art) administration into the skin of mice produced an intense accumulation of polymorphonuclear cells, largely neutrophils. In contrast, fMLPP (HK-X) has no significant effect when administered alone. However, in connective tissues HK-X administered simultaneously with an equal amount of fMLP totally prevented the accumulation and diapedesis of polymorphonuclear cells into interstitial spaces while administration of HK-X alone showed no inflammatory recruitment of polymorphonuclear cells or other inflammatory cells in the connective tissues. It can also be concluded that HK-X blocks the earliest events triggered by fMLP in the inflammation cascade, that is, recruitment of inflammatory cells.

18. It is my understanding that the Examiner contends that, based on Kermode et al, "one would expect that fMLPP, alone, would be at least as "pro-inflammatory" as fMLP." The above described experiments show the opposite result, that is, fMLPP is anti-inflammatory and also inhibits the pro-inflammatory effects of fMLP.

19. I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true;

Declaration of James Clagett, Ph.D.
Ser. No. 09/189,130
Page 8 of 8

8

and, further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Codes, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

11-21-00
Date




James Clagett

FIGURE 1

HK-X Prevents fMLP Induced Neutrophil Migration



Figures 2

Micrographs to illustrate the treatment reaction in skins of mice.

A). fMLP treated mouse skin shows the influx of the neutrophils in the surrounding areas of the blood vessel (BV). The neutrophils are located in the connective tissue area (Arrows).

B). A fMLP and HK-X treated mouse skin shows no cellular infiltration at the surrounding areas of the blood vessel (BV). Occasionally neutrophils are seen (Arrows).

C). HK-X treated mouse skin shows no changes in the connective tissues and the surrounding area of the blood vessel (BV).










Figure 13

A). A HK-X treated mouse skin shows the normal appearances of the subcutaneous tissue. The blood vessels (BV) show no neutrophils. Only occasionally the observation of neutrophil present (Arrow).


B). A vehicle control mouse skin shows no sign of the effect of vehicle solution injection.

C). A HK-X and fMLP treated mouse skin shows very little change in the histology except a few neutrophils are attaching on the endothelial cell surface (Arrow heads). Neutrophils are seen but in very little number as compared to the fMLP treated skin tissue (see Fig. D)

D). A fMLP treated mouse skin shows the abundant number of neutrophils present in the surrounding area of the blood vessel (BV). Many neutrophils are also seen attached to the surface of endothelial cells.



JAMES A. CLAGETT, Ph.D.
5615 139th Avenue SE
Snohomish, Washington 98290



Telephone
Residence: (360) 568-3988
Business: (360) 568-3988

EXPERIENCE:

CLAGETT CONSULTING
5615 139TH AVE SE
SNOHOMISH, WASHINGTON, 98290

Founder and Partner: January 1, 1997 to present.

Providing quality expertise to the biotechnology and pharmaceutical communities. Lead an assembled group of individuals with experience in regulatory affairs, preclinical and clinical studies. The team has worked together for an average of 3 years and was instrumental in bringing products from research through production for GenSci Regeneration Laboratories Inc., Irvine, CA. Current list of clients includes Histatek, LLC, San Francisco, CA and BioTherapeutic Computers, Seattle, WA.

BIOCOLL LABORATORIES INC.
562 1ST AVENUE SOUTH
SEATTLE, WASHINGTON, 98104

Vice President and Scientific Director: January 1993 to August 1, 1997.

Responsible for implementing and maintaining the company's overall scientific plan. Initiated the development of the company's first product, the Tissue Bone Matrix sponge and the DynaGraft family of products and human banked tissue. Supervised the writing and implementation of the Standard Operating procedures for the TBM sponge. Initiated the scientific research of new products and managed numerous consultants supporting the company's scientific strategic plan. Carried the company from conceptual design of products through introduction to the marketplace including continuous interaction with FDA.

DENTAL DIAGNOSTIC SERVICES
2000 116TH AVENUE NE.
BELLEVUE, WASHINGTON 98004

Founder and Officer: December 1992 through January 1, 1993

Responsible for taking the company from no revenues to profitability in one year. Company had over 450 clients using the sterility testing services of Dental Diagnostic Services.

THE DENTAL RESOURCE INC.
2000 116TH AVENUE NE,
BELLEVUE, WASHINGTON 98004

Consultant: October 1990 to December 1992

Responsible for developing product concept for novel water filtration device for dental offices. Participated heavily in selling devices to dental practitioners.

ULTRA DIAGNOSTICS CORPORATION
4526 11th Avenue NE, Seattle, Washington, 98108

Chief Executive Officer, President, and Scientific Director: December 1988 to December 1989.

Responsible for implementing and maintaining the company's overall business plans. Maintained on-going communications with investors. Supervised the recruitment of research and development personnel and management finance, administration, marketing and regulatory functions. Responsible of business development where a contract of \$500,000 for research and development of new reagents was concluded. Responsible for continued venture funding of approximately \$900,000. Maintained ongoing contact with key administrative and upper management individuals of the biotechnology and academic sectors in the Northwest.

President and Scientific Director: December 1984 to December 1988. Implemented the company's research and development plans and assisted the Chief Executive Officer with business development functions where research and development contracts worth \$500,000 were secured. Coordinated the use of the Scientific Advisory Board and recruited and hired key scientific personnel. Assisted the Chief Executive Officer in capital formation from "seed" through venture financing. The capital raised was in excess of \$4 million.

**UNIVERSITY OF WASHINGTON
Schools of Medicine and Dentistry
Seattle, Washington, 98105**

Affiliate Professor: 1987 to 1989.

Taught a course in Immunology and Immunopathology to Graduate Dental Students.

Professor of Periodontics and Microbiology and Immunology: 1983 to 1987

Associate Professor of Periodontics and Microbiology and Immunology: 1978 to 1983.

Research Assistant Professor of Periodontics and Microbiology and Immunology: 1973 to 1978

Taught graduate and undergraduate Immunology and Immunopathology to Microbiology and Immunology majors as well as participated in team teaching to dental and medical students. Received well above average reviews of teaching skills by students. Maintained a research laboratory with technicians, 2 doctoral and 6 masters' students. Obtained research support averaging \$150,000 per year and served on the Faculty Senate and the Human Subjects Review Committee. Served as a member on 15 doctoral and masters thesis committees.

PROFESSIONAL ASSOCIATIONS:

Editorial Board of Journal of Dental Research
Biomedical Research Support Grant Committee
American Association of immunologists
Sigma Xi

EDUCATION:

Scripps Clinic and Research Foundation	Postdoctoral Fellowship	1973
University of Nebraska	Ph.D.	1970
DePauw University	BS	1964

PUBLICATIONS

Clagett, J. A., Engelhard, W. E. The immunosuppressive activity of the 3-a1 fraction of papain. *J. Bacteriol.* 93:19492, 1967.

Peter, H. H., **Clagett, J. A.,** Feldman, J. D., and Weigle, W. O. Rabbit antiserum to brain associated thymus antigens of mouse and rat. I. Demonstration of antibodies cross-reacting to T cells of both species. *J. Immunol.* 110:1077, 1973.

Clagett, J. A., Peter, H. H., Feldman, J. D., and Weigle, W. O. "Rabbit antiserum to brain associated thymus antigens of mouse and rat. II. Detection of species specific and theta related determinants. *J. Immunol.* 110:1085, 1973.

Clagett, J. A., Tokuda, S., Engelhard, W. E. Chymopapain C., an immunosuppressive protease: I. Purification and characterization. *Proc. Soc. Exp. Biol. Med.*, 145:1250, 1974.

Clagett, J. A., Tokuda, S., and Engelhard, W. E. Chymopapain C., an immunosuppressive protease: II. Effect on the primary and secondary response in mice. *J. Immunol.* 112:1669, 1974.

Clagett, J. A., and Weigle, W. O. Role of T- and B-lymphocytes in the termination of unresponsiveness to autologous thyroglobulin. *J. Exp. Med.* 139:643, 1974.

Clagett, J. A., Wilson, C. B., and Weigle, W. O. Interstitial immune complex in mice: The role of autoantibody to thyroglobulin. *J. Exp. Med.* 140:1439, 1974.

Huang, L. Y., Stern, I. B., **Clagett, J. A.,** and Chi, E. Y. Two polypeptide chain constituents of the major protein of the cornified layer of newborn rat epidermis. *Biochemistry* 14:3573, 1975.

Engel, L. D., Van Epps, D., and **Clagett, J. A.** In vivo and in vitro studies on possible pathogenic mechanisms of *Actinomyces viscosus*. *Infection and Immunity* 14:548, 1976.

Storb, U., Hager, L., Putnam, D., Buck, L., Marvin, S., Farin, F., and **Clagett, J.** Sequences related to immunoglobulin kappa chain messenger RNA T cells. *Proc. Nat'l. Acad. Sci. U.S.A.* 73:2467, 1976

Dale, B., Stern, I. B., and **Clagett, J. A.** Initial characterization of the proteins of keratinized epithelium of rat oral mucosa. *Arch. Oral Biol* 22:75, 1977.

Putnam, D., Storb, U., and **Clagett, J.** Synthesis of kappa chains from thymus RNA by cell free translation. Proceedings of UC and ICN Meeting, March, 1977. In Regulatory Genetics of the Immune System ICN-UCLA Symposia on Molecular and Cellular Biology. 6:71, 1977.

Press, O. W., Rosse, C. and **Clagett, J. A.** The distribution of short-lived and long-lived T-, B- and null-lymphocytes mouse bone marrow, thymus, lymph node and spleen. *Cellular Immunol.* 33:114-124, 1977.

Morton, T., **Clagett, J. A.,** and Yavorsky, D. The role of immune complexes in chronic periapical lesions. *J. Endodontics* 3:261, 1977.

- Engel, L. D., **Clagett, J. A.**, Page, R. C., and Williams, B. Mitogenic activity of Actinomyces viscosus: I. Effects of murine B- and T-lymphocytes and partial characterization. *J. Immunol.* 118:1466, 1977.
- Press, O. W., Rosse, C., and **Clagett, J. A.** Phytohemagglutinin-induced differentiation and blastogenesis of precursor T-cells from mouse bone marrow. *J Exp. Med.* 146~735 1977.
- Press, O. W., Rosse, C., and **Clagett, J. A.** Anti- θ brain antisera bind primarily to non-T cells in mouse bone marrow. *Proc. Soc. Exp. Biol. Med* 156:485 ,1977.
- Clagett, J. A.**, and Page, R. C. Insoluble immune complexes and periodontal diseases in man and the dog. *Arch. Oral Biol.* 23:153, 1978.
- Clagett, J. A.**, and Engel, L. D. Polyclonal activation: A form of primitive immunity and its possible role in chronic inflammatory diseases. *Develop. and Comp. Immunol.* 2:235, 1978.
- Page, R. C., Engel, L. D., **Clagett, J. A.**, and Narayanan, A. S. Chronic inflammatory gingival and periodontal disease. *J. Am. Med. Assoc.* 240:545, 1978.
- Rosse, C., Cole, S. B., Appleton, C., Press, O. W., and **Clagett, J. A.** The relative importance of the bone marrow in the production and dissemination of B-lymphocytes. *Cell Immunol.* 37:254 , 1978.
- Sims, T., **Clagett, J. A.**, and Page, R. C. Effects of cell concentration and exogenous prostaglandin on the interaction and responsiveness of human peripheral blood leukocytes. *Clin. Immunol. and Immuno path.* 12: 150, 1979.
- Page, R. C., **Clagett, J. A.**, Engel, L. D., and Sims, T. Effects of prostaglandin on the antigen and mitogen-driven responses of peripheral blood lymphocytes from patients with adult and juvenile periodontitis. *Clin. Immunol. and Immunopath.* 11:77 , 1978.
- Torbinejab, M., **Clagett, J.**, and Engel, D. A cat model for the evaluation of mechanisms of bone resorption: induction of bone loss by stimulated immune complexes and inhibition by Indomethacin. *Calcif. Tissue Res.* 29:207, 1979.
- Putnam, D., **Clagett, J.**, and Storb, U. Immunoglobulin synthesis by T cells: quantitative and qualitative aspects. *J Immunol.* 124:920, 1980.
- Storb, U., Near, R., **Clagett, J.** Immunoglobulin messenger RNA's of T lymphocytes. *Molecular Immunology* 17:474, 1980.
- Engel, L. D., Chi, E., and **Clagett, J. A.** Mitogenic activity of Actinomyces viscosus. II. Induction of DNA and immunoglobulin synthesis in rabbit B lymphocytes. *Dev. Comp. Immunol* 4:515 1980.
- Engel, L. D., Schroder, H. E., Gay, R. G., and **Clagett, J. A.** Fine structure of cultured human gingival fibroblasts and demonstration of simultaneous synthesis of types I and III collagen. *Arch. Oral Biol.* 25:283 , 1980.
- Clagett, J. A.**, Engel, L. D., and Chi, E. In vitro expression of immunoglobulin M and G subclasses by murine B lymphocytes in response to a polyclonal activator from Actinomyces viscosus. *Infect. Immunity* 1980.

Korotzer, T. I., **Clagett, J. A.**, Kolb W. P., and Page, R. C. Complement dependent induction of DNA synthesis and proliferation of human diploid fibroblasts *J. Cell Physiol.* 105:503, 1980.

Landreth, K. S, McCoy, K. L., **Clagett, J. A.**, Bollum, F. J., and Rosse, C. Deficiency in terminal deoxynucleotidyl transferase-positive cells associated with severe autoimmune disease. *Nature* 290~409 1981.

Engel, L. D., Clark, E. A., Held, L., Kimball, H., and **Clagett, J. A.** Immune responsiveness of SM/J mice. Cellular characteristics and genetic analysis of hyperresponsiveness to B cell mitogens. *J Exp. Med.* 154:726,1981.

Landreth, D. S., Ross, C., and **Clagett, J. A.** The myelogenous phase in the production and maturation of primary B lymphocytes in the mouse. *J. Immunology* 127:2027, 1981.

McCoy, K. L., **Clagett, J. A.**, and Chi, E. Abnormal maturation of mono-nuclear phagocytes associated with autoimmunity in motheaten mice. *Oral Immunogenetics and Tissue Transplantation*, G. R. Riviere and W. H. Hildemann, eds. Elsevier/North Holland, 1982.

McCoy, K. L., Chi, E., Engel, D., and **Clagett, J.** Abnormal in vitro proliferation of splenic mononuclear phagocytes from autoimmune motheaten mice. *J. Immunology* 128:1797,1982.

McCoy, K. L., and **Clagett, J. A.** Accelerated rate of mononuclear phagocyte maturation in autoimmune motheaten mice. *Fed. Proc.* 41:559,1982.

Engel, D., Clark, E. A., Chi, E., and **Clagett, J.** *Genetic aspects of responsiveness to B cell mitogens in mice.* *Oral Immunogenetics and Tissue Transplantation*, G. R. Riviere, ed. Elsevier/North Holland, 1982.

McCoy, K. L., **Clagett, J. A.**, Engel, L. D., and Chi, E. Accelerated rate of mononuclear phagocyte production in vitro by splenocytes from autoimmune motheaten mice. *Am J. Pathol.* 112:18 1983.

Clagett, J. A., and Engel, L. D. *Polyclonal B-cell activation in response to Actinomyces viscosus - its nature and genetics.* In Biological, Biochemical and Biomedical Aspects of Actinomycetes. Edited by L. Ortiz-Ortiz, L. F. Bojalil and V. Yakoleff. Academic Press, Orlando, pp.61-71,1984.

McCoy, K. L., Nielson, K., and **Clagett, J.** Spontaneous Production of Colony-Stimulating Activity by Splenic Mac-I Antigen-Positive Cells from Autoimmune Motheaten Mice. *J Immunology* :272, 1984.

Engel, D., Monzingo, S., Rabinovitch, P., **Clagett, J.**, and Stone, R. Mitogen-induced hyperproliferation response of peripheral blood mononuclear cells from patients with severe generalized periodontitis: lack of correlation with proportions of T-cells and T-cell subsets.. *Immunopathol Clin. Immunol* 30:374, 1984.

McAnulty, K., Stone, R., Hastings, G., **Clagett, J.**, and Engel, D. Immunoregulation in severe generalized periodontitis. *Clin. Immunol. Immunopathol* 34:84 1985.

ABSTRACTS

Clagett, J. A. Cellular collaboration and specificity in murine thyroiditis. *Fed. Proc.* 31:743, 1972.

Peter, H. H., **Clagett, J. A.**, Wilson, C. B., and Feldman, J. D. Rabbit antisera to brain associated thymus antigens of mouse and rat. *German Soc. Immunol.*, 1972.

Engel, L. D., **Clagett, J. A.**, and Page, R. C. A possible role for bacterial mitogens in chronic inflammatory periodontal disease. *J Periodontol.*, 1975.

Engel, L. D., **Clagett, J. A.**, and Page, R. C. Response of mouse spleen cells to Actinomyces viscosus antigens. *FASEB*, 1975.

Clagett, J. A., Booth, R., and Page, R. C. Role of immune complexes in human periodontitis. *IADR*, 1976.

Suzuki, J. B., **Clagett, J. A.** Sims, T., and Page R. C. Analysis of the cell-cell interactions in the blastogenic response of lymphocytes of periodontal disease patients. *IADR*, 1979.

Putnam D. L. **Clagett, J. A.** Different proliferation and immunoglobulin responses of B cells to four mitogens. *FASEB*, 1980.

McCoy, K. Rosse, C., and **Clagett J.** The cellular composition and B cell mitogen responsiveness of lymphoid cells of motheaten mice (me/me). *FASEB*, 1980.

Landreth, K. S., Rosse, C., and **Clagett, J.** Evidence for a lineage relationship between cytoplasmic micron+ micron-, and surface micron+ B lymphocytes in adult murine bone marrow. *FASEB*, 1980.

Clagett, J. A., klimpel, K., Klimpel, G., and McCoy, K. Ia-negative murine B cells: Proliferative and differentiative capacity in vitro. *IADR*, 1981.

Engel, D., Kimbalil H., Clark, E., and **Clagett, J.** Genetic control of high responsiveness to B-cell mitogens. *IADR*, 1981.

McCoy, K., Rosse, C., and Clagett, J. Abnormal proliferation of splenic macrophages from autoimmune motheaten mice. *FASEB*, 1981.

Engel, D., Monzingo S., Rabinovitch, P., **Clagett, J.** Mitogen-induced hyperproliferation response of lymphocytes from patients with severe generalized periodontitis: Lack of correlation with proportions of T-cell subsets. *FASEB* 1982.

Windsor, N., **Clagett, J. A** Novel effector system: Bone marrow transformation activity produced by a T:T hybridoma. *FASEB*, 1984.

Beneveniste, R. E., Stromberg, K., Arthur, L. O., Giddens, W. E. Jr., Ochs, H. D., Morton, W. R., and **Clagett, J. A.** Transmission of Retroperitoneal Fibromatosis (RF) by SAIDSD/Washington retrovirus. *International Academy of Pathology*, 1985.